


# AMENDMENT TO THE CLAIMS

Please amend the claims as follows:

1. (currently amended) A compound of formula I below, and physiologically acceptable salts, comprising:



wherein,

the "A" ring atoms of compound formula I comprise carbon and 0 to 2 nitrogen heteroatoms;

Ar is an aromatic ring, an aromatic ring comprising at least one substituent group, a heteroaromatic ring, a heteroaromatic ring comprising 1 to 5 substituent groups, a heterocyclic ring or a heterocyclic ring comprising at least one substituent group;

R ~~comprises~~ is selected from H, OH, OCH<sub>3</sub>, alkoxy, OCH<sub>2</sub>CH<sub>2</sub>OH, alcohol, NH<sub>2</sub>, PO<sub>3</sub>H, OPO<sub>3</sub>H, OSO<sub>3</sub>H, halogen, C(halogen)<sub>3</sub>, SE<sub>1</sub>, OE<sub>1</sub> ~~or~~ and NE<sub>1</sub>E<sub>2</sub>,

E<sub>1</sub> and E<sub>2</sub> are each independently H or alkyl;

R' ~~comprises~~ is selected from H, OH, alkoxy, OCH<sub>2</sub>CH<sub>2</sub>OH, alcohol, NH<sub>2</sub>, PO<sub>3</sub>H, OPO<sub>3</sub>H, OSO<sub>3</sub>H, halogen, C(halogen)<sub>3</sub>, SE<sub>1</sub>, OE<sub>1</sub> ~~or~~ and NE<sub>1</sub>E<sub>2</sub>,

E<sub>1</sub> and E<sub>2</sub> are each independently H or alkyl;

~~R'' and R''' each independently comprises~~ is Y-D<sub>1</sub>-D<sub>2</sub>-T<sub>2</sub>, H, ~~halogen, alkyl, alkoxy or a substituent group,~~

Y is optionally present and if present ~~comprises~~ is selected from O, S, NH, N-alkyl, C=CH, C≡C, CH<sub>2</sub>, ~~CH(CH<sub>3</sub>)~~ CH(CH<sub>3</sub>), C(CH<sub>3</sub>)<sub>2</sub>, a carbocyclic ring having 4

to 6 ring members ~~or~~ and a heterocyclic ring having 4 to 6 ring members with 1 or 2 heteroatoms,

D<sub>1</sub> is optionally present and if present ~~comprises~~ is alkyl,

D<sub>2</sub> ~~comprises~~ is selected from H, alkyl, NH, N-alkyl, O-alkyl, S-alkyl, a carbocyclic ring, a bicyclic ring, a tricyclic ring, an aromatic ring ~~or~~ and a heteroaromatic ring,

T<sub>2</sub> is optionally present and if present ~~comprises~~ is selected from an aromatic ring, a substituted aromatic ring, a heteroaromatic ring, a substituted heteroaromatic ring, a heterocyclic ring, a substituted heterocyclic ring, H, OH, halogen, ~~or~~ and a substituent group;

R''' and R'''' are each independently selected from H, halogen, alkyl, alkoxy and a substituent group;

with the proviso that,

when Ar is 4-isopropyl pyridine or 4-isopropenyl pyridine, R''' is hydrogen, and R'''' is hydrogen, then R'' can not be a straight or branched saturated alkyl having 1 to 20 carbon atoms;

when Ar is 4-isopropyl toluene or 4-isopropenyl toluene, and both R''' and R'''' are hydrogen, R'' can not be a straight or branched saturated alkyl having 1 to 20 carbon atoms;

when R'' is C(CH<sub>3</sub>)<sub>2</sub>(CH<sub>2</sub>)<sub>5</sub>CH<sub>3</sub>, R<sub>2</sub> and R<sub>4</sub> are methyl, then R' and R'' can not be H, OH or OCH<sub>3</sub>.

2. cancelled

3. (currently amended) The compound of claim 1 wherein:

R''' ~~comprises~~ is selected from H, halogen, C(halogen)<sub>3</sub>, lower alkyl ~~or~~ and alkoxy;

R'''' ~~comprises~~ is selected from H, halogen, C(halogen)<sub>3</sub>, lower alkyl ~~or~~ and alkoxy; and

R'' ~~comprises~~ is selected from -Y-D<sub>1</sub>-D<sub>2</sub>-T<sub>2</sub>,

Y ~~comprises~~ is selected from  $C(CH_3)_2$ ,  $CH_2$  ~~or~~ and  $CH(CH_3)$ ,

$D_1$  is optionally present and if present ~~comprises~~ is alkyl,

$D_2$  ~~comprises~~ is selected from H, an alkyl, NH, N-alkyl, O-alkyl, S-alkyl, a carbocyclic ring, a bicyclic ring, a tricyclic ring, an aromatic ring ~~or~~ and a heteroaromatic ring,

$T_2$  is optionally present and if present ~~comprises~~ is selected from an aromatic ring, a heteroaromatic ring, a heterocyclic ring, H, OH, halogen ~~or~~ and a substituent group.

4. (currently amended) The compound of claim 1 wherein:

$R'''$  ~~comprises~~ is selected from H, halogen,  $C(halogen)_3$ , lower alkyl ~~or~~ and alkoxy;

$R''''$  ~~comprises~~ is selected from H, halogen,  $C(halogen)_3$ , lower alkyl ~~or~~ and alkoxy; and

$R''$  ~~comprises~~ is  $-Y-D_1-D_2-T_2$ ,

Y ~~comprises~~ is selected from O, NH ~~or~~ and N-alkyl,

$D_1$  is optionally present and if present ~~comprises~~ is alkyl,

$D_2$  ~~comprises~~ is selected from H, an alkyl, NH, N-alkyl, O-alkyl, S-alkyl, a carbocyclic ring, a bicyclic ring, a tricyclic ring, an aromatic ring ~~or~~ and a heteroaromatic ring,

$T_2$  is optionally present and if present ~~comprises~~ is selected from an aromatic ring, a heteroaromatic ring, a heterocyclic ring, H, OH, halogen ~~or~~ and a substituent group.

5. (currently amended) The compound of claim 1 wherein:

$R'''$  ~~comprises~~ is selected from H, halogen,  $C(halogen)_3$ , lower alkyl ~~or~~ and alkoxy;

$R''''$  ~~comprises~~ is selected from H, halogen,  $C(halogen)_3$ , lower alkyl ~~or~~ and alkoxy; and

$R''$  ~~comprises~~ is  $-Y-D_1-D_2-T_2$ ,

Y is optionally present and if present ~~comprises~~ is selected from  $C=CH$  ~~or~~ and  $C\equiv C$ ,

D<sub>1</sub> is optionally present and if present ~~comprises~~ is alkyl,

D<sub>2</sub> ~~comprises~~ is selected from H, alkyl, NH, N-alkyl, O-alkyl, S-alkyl, a carbocyclic ring, a bicyclic ring, a tricyclic ring, an aromatic ring ~~or~~ and a heteroaromatic ring,

T<sub>2</sub> is optionally present and if present ~~comprises~~ is selected from an aromatic ring, a heteroaromatic ring, a heterocyclic ring, H, OH, halogen ~~or~~ and a substituent group.

6. (currently amended) The compound of claim 1 wherein:

R''' ~~comprises~~ is selected from H, halogen, C(halogen)<sub>3</sub>, lower alkyl ~~or~~ and alkoxy;

R''' ~~comprises~~ is selected from H, halogen, C(halogen)<sub>3</sub>, lower alkyl ~~or~~ and alkoxy; and

R'' ~~comprises~~ is -Y-D<sub>1</sub>-D<sub>2</sub>-T<sub>2</sub>,

Y ~~comprises~~ is optionally present and if present is selected from 0 to 1 of a carbocyclic ring having 4 to 6 ring members ~~or~~ and a heterocyclic ring having 4 to 6 ring members with 1 or 2 heteroatoms.



D<sub>1</sub> is optionally present and if present ~~comprises~~ is alkyl,

D<sub>2</sub> ~~comprises~~ is selected from H, alkyl, NH, N-alkyl, O-alkyl, S-alkyl, a carbocyclic ring, a bicyclic ring, a tricyclic ring, an aromatic ring ~~or~~ and a heteroaromatic ring,

T<sub>2</sub> is optionally present and if present ~~comprises~~ is selected from an aromatic ring, a heteroaromatic ring, a heterocyclic ring, H, OH, halogen ~~or~~ and a substituent group.

7. (currently amended) The compound of claim 1 wherein Ar ~~comprises~~ is selected from an aromatic ring having 5 or 6 ring members ~~or~~ and a heteroaromatic ring having 5 or 6 ring members.

8. (currently amended) The compound of claim 1 wherein Ar comprises is selected from one of the structures:




and,

the Ar aromatic ring structure comprises 0 to 3 heteroatoms as ring members;

R1, R2, R3, R4 and R5 are each independently comprise selected from H, OH, NH<sub>2</sub>, halogen, N<sub>3</sub>, NO<sub>2</sub>, NCS, C(halogen)<sub>3</sub>, CHO, OAc, OCH<sub>3</sub>, OC<sub>2</sub>H<sub>5</sub>, CH<sub>2</sub>OH, CH<sub>2</sub>CH<sub>2</sub>OH, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OH, CN, C(=O)CH<sub>3</sub>, COOH, COOCH<sub>3</sub>, COOC<sub>2</sub>H<sub>5</sub>, COOCH(CH<sub>3</sub>)<sub>2</sub>, NHCOCH<sub>3</sub>, SCH<sub>3</sub>, SC<sub>2</sub>H<sub>5</sub>, NHCH<sub>3</sub>, CH<sub>2</sub>NH<sub>2</sub>, CH<sub>3</sub>, C<sub>2</sub>H<sub>5</sub>, C<sub>3</sub>H<sub>7</sub>, C<sub>2</sub>H<sub>3</sub>, ethynyl, alkoxy, alkylmercapto, alkylamino, di-alkylamino, alkylsulfinyl, alkylsulfonyl, or methylene dioxy or and a substituent group.


9. (currently amended) The compound of claim 1 wherein Ar comprises is selected from 1-, 2- or 3-pyrrolidinyl, 1-, 2-, 3- or 4-piperidinyl, 1-, 2- or 3-morpholinyl, 1-, 2- or 3-thiomorpholinyl, 1-, 2- or 3- azetidiny, 1-, or 2-piperazinyl, 2- or 3-tetrahydrofuranyl; or any above group substituted on any available ring carbon thereof by alkyl; or any above group unsubstituted on one or more nitrogen atoms, or any above group substituted on one or more nitrogen atoms independently by an alkyl, benzyl, lower-alkoxybenzyl or benzhydryl group; adamantyl; a carbocyclic ring, a substituted carbocyclic ring, a heteroaromatic ring, a substituted heteroaromatic ring, a heterocyclic ring, a substituted heterocyclic ring, a bicyclic ring, a substituted bicyclic ring, a heterobicyclic ring, a substituted heterobicyclic ring, a polycyclic ring, a substituted polycyclic ring, a heteropolycyclic ring or a substituted heteropolycyclic ring.

10. (currently amended) The compound of claim 1 wherein Ar comprises is selected from:



G comprises is selected from H, OH, NH<sub>2</sub>, halogen, N<sub>3</sub>, NO<sub>2</sub>, NCS, CF<sub>3</sub>, CHO, OAc, OCH<sub>3</sub>, OC<sub>2</sub>H<sub>5</sub>, CH<sub>2</sub>OH, CH<sub>2</sub>CH<sub>2</sub>OH, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OH, CN, C(=O)CH<sub>3</sub>, COOH, COOCH<sub>3</sub>, COOC<sub>2</sub>H<sub>5</sub>, COOCH(CH<sub>3</sub>)<sub>2</sub>, NHCOCH<sub>3</sub>, SCH<sub>3</sub>, SC<sub>2</sub>H<sub>5</sub>, NHCH<sub>3</sub>, CH<sub>2</sub>NH<sub>2</sub>, CH<sub>3</sub>, C<sub>2</sub>H<sub>5</sub>, C<sub>3</sub>H<sub>7</sub>, C<sub>2</sub>H<sub>3</sub>, ethynyl, alkoxy, alkylmercapto, alkylamino, di-alkylamino, alkylsulfinyl, alkylsulfonyl or and methylene dioxy.

11. (currently amended) A pharmaceutical preparation comprising a therapeutically effective amount of at least one compound of formula I below, and physiologically acceptable salts thereof:



wherein,

the "A" ring atoms of compound formula I comprise carbon and 0 to 2 nitrogen heteroatoms;

Ar is an aromatic ring, an aromatic ring comprising at least one substituent group,

a heteroaromatic ring, a heteroaromatic ring comprising 1 to 5 substituent groups, a heterocyclic ring or a heterocyclic ring comprising at least one substituent group;

R ~~comprises~~ is selected from H, OH, OCH<sub>3</sub>, alkoxy, OCH<sub>2</sub>CH<sub>2</sub>OH, alcohol, NH<sub>2</sub>, PO<sub>3</sub>H, OPO<sub>3</sub>H, OSO<sub>3</sub>H, halogen, C(halogen)<sub>3</sub>, SE<sub>1</sub>, OE<sub>1</sub> ~~or~~ and NE<sub>1</sub>E<sub>2</sub>,

E<sub>1</sub> and E<sub>2</sub> are each independently H or alkyl;

R' ~~comprises~~ is selected from H, OH, alkoxy, OCH<sub>2</sub>CH<sub>2</sub>OH, alcohol, NH<sub>2</sub>, PO<sub>3</sub>H, OPO<sub>3</sub>H, OSO<sub>3</sub>H, halogen, C(halogen)<sub>3</sub>, SE<sub>1</sub>, OE<sub>1</sub> ~~or~~ and NE<sub>1</sub>E<sub>2</sub>,

E<sub>1</sub> and E<sub>2</sub> are each independently H or alkyl;

R''~~[[,]]~~ R''' and R'''' each independently comprises is Y-D<sub>1</sub>-D<sub>2</sub>-T<sub>2</sub>, H, halogen, ~~alkyl, alkoxy or a substituent group,~~

Y is optionally present and if present ~~comprises~~ is selected from O, S, NH, N-alkyl, C=CH, C≡C, CH<sub>2</sub>, CH(CH<sub>3</sub>), C(CH<sub>3</sub>)<sub>2</sub>, a carbocyclic ring having 4 to 6 ring members ~~or~~ and a heterocyclic ring having 4 to 6 ring members with 1 or 2 heteroatoms,

D<sub>1</sub> is optionally present and if present ~~comprises~~ is alkyl,

D<sub>2</sub> ~~comprises~~ is selected from H, alkyl, NH, N-alkyl, O-alkyl, S-alkyl, a carbocyclic ring, a bicyclic ring, a tricyclic ring, an aromatic ring ~~or~~ and a heteroaromatic ring,

T<sub>2</sub> is optionally present and if present ~~comprises~~ is selected from an aromatic ring, a substituted aromatic ring, a heteroaromatic ring, a substituted heteroaromatic ring, a heterocyclic ring, a substituted heterocyclic ring, H, OH, halogen, ~~or~~ and a substituent group;

R''' and R'''' are each independently selected from H, halogen, alkyl, alkoxy and a substituent group;

with the proviso that,

when Ar is 4-isopropyl pyridine or 4-isopropenyl pyridine, R''' is hydrogen, and R'''' is hydrogen, then R'' can not be a straight or branched saturated alkyl having 1 to 20 carbon atoms;

when Ar is 4-isopropyl toluene or 4-isopropenyl toluene, and both R''' and R'''' are hydrogen, R'' can not be a straight or branched saturated alkyl having 1 to 20 carbon atoms;

when R'' is C(CH<sub>3</sub>)<sub>2</sub>(CH<sub>2</sub>)<sub>5</sub>CH<sub>3</sub>, R<sub>2</sub> and R<sub>4</sub> are methyl, then R' and R'' can not be H, OH or OCH<sub>3</sub>.

12. cancelled

13. (currently amended) The pharmaceutical preparation of claim 11, wherein:

R''' comprises is selected from H, halogen, C(halogen)<sub>3</sub>, lower alkyl ~~or~~ and alkoxy;

R'''' comprises is selected from H, halogen, C(halogen)<sub>3</sub>, lower alkyl ~~or~~ and alkoxy; and

R'' comprises is -Y-D<sub>1</sub>-D<sub>2</sub>-T<sub>2</sub>,

~~Y~~ comprises Y is optionally present and if present comprises is selected from O, S, NH, N-alkyl, C=CH, C≡C, CH<sub>2</sub>, ~~CH(CH<sub>3</sub>)~~ CH(CH<sub>3</sub>), C(CH<sub>3</sub>)<sub>2</sub>, a carbocyclic ring having 4 to 6 ring members ~~or~~ and a heterocyclic ring having 4 to 6 ring members with 1 or 2 heteroatoms,

D<sub>1</sub> is optionally present and if present comprises is alkyl,


D<sub>2</sub> comprises is selected from H, ~~an~~ alkyl, NH, N-alkyl, O-alkyl, S-alkyl, a carbocyclic ring, a bicyclic ring, a tricyclic ring, an aromatic ring ~~or~~ and a heteroaromatic ring,

T<sub>2</sub> is optionally present and if present comprises is selected from an aromatic ring, a heteroaromatic ring, a heterocyclic ring, H, OH, halogen ~~or~~ and a substituent group.

14. (currently amended) A method of stimulating a cannabinoid receptor in an



individual or animal comprising administering to the individual or animal a therapeutically effective amount of ~~a therapeutically effective amount~~ of at least one compound of formula I below, and physiologically acceptable salts thereof:



wherein,

the "A" ring atoms of compound formula I comprise carbon and 0 to 2 nitrogen heteroatoms;

Ar is an aromatic ring, an aromatic ring comprising at least one substituent group, a heteroaromatic ring, a heteroaromatic ring comprising 1 to 5 substituent groups, a heterocyclic ring or a heterocyclic ring comprising at least one substituent group;

R ~~comprises~~ is selected from H, OH, OCH<sub>3</sub>, alkoxy, OCH<sub>2</sub>CH<sub>2</sub>OH, alcohol, NH<sub>2</sub>, PO<sub>3</sub>H, OPO<sub>3</sub>H, OSO<sub>3</sub>H, halogen, C(halogen)<sub>3</sub>, SE<sub>1</sub>, OE<sub>1</sub> ~~or~~ and NE<sub>1</sub>E<sub>2</sub>,

E<sub>1</sub> and E<sub>2</sub> are each independently H or alkyl;

R' ~~comprises~~ is selected from H, OH, alkoxy, OCH<sub>2</sub>CH<sub>2</sub>OH, alcohol, NH<sub>2</sub>, PO<sub>3</sub>H, OPO<sub>3</sub>H, OSO<sub>3</sub>H, halogen, C(halogen)<sub>3</sub>, SE<sub>1</sub>, OE<sub>1</sub> ~~or~~ and NE<sub>1</sub>E<sub>2</sub>,

E<sub>1</sub> and E<sub>2</sub> are each independently H or alkyl;

~~R''[,], R''' and R''' each independently comprises~~ is Y-D<sub>1</sub>-D<sub>2</sub>-T<sub>2</sub>, H, ~~halogen, alkyl, alkoxy or a substituent group,~~

Y is optionally present and if present ~~comprises~~ is selected from O, S, NH, N-alkyl, C=CH, C≡C, CH<sub>2</sub>, ~~CH(CH<sub>3</sub>)~~ CH(CH<sub>3</sub>), C(CH<sub>3</sub>)<sub>2</sub>, a carbocyclic ring having 4 to 6 ring members ~~or~~ and a heterocyclic ring having 4 to 6 ring members with 1 or 2 heteroatoms,

D<sub>1</sub> is optionally present and if present ~~comprises~~ is alkyl;

D<sub>2</sub> ~~comprises~~ is selected from H, alkyl, NH, N-alkyl, O-alkyl, S-alkyl, a carbocyclic ring, a bicyclic ring, a tricyclic ring, an aromatic ring ~~or~~ and a heteroaromatic ring,

T<sub>2</sub> is optionally present and if present ~~comprises~~ is selected from an aromatic ring, a substituted aromatic ring, a heteroaromatic ring, a substituted heteroaromatic ring, a heterocyclic ring, a substituted heterocyclic ring, H, OH, halogen, ~~or~~ and a substituent group;

R''' and R'''' are each independently selected from H, halogen, alkyl, alkoxy and a substituent group;

with the proviso that,

when Ar is 4-isopropyl pyridine or 4-isopropenyl pyridine, R''' is hydrogen, and R'''' is hydrogen, then R'' can not be a straight or branched saturated alkyl having 1 to 20 carbon atoms;

when Ar is 4-isopropyl toluene or 4-isopropenyl toluene, and both R''' and R'''' are hydrogen, R'' can not be a straight or branched saturated alkyl having 1 to 20 carbon atoms;

when R'' is C(CH<sub>3</sub>)<sub>2</sub>(CH<sub>2</sub>)<sub>5</sub>CH<sub>3</sub>, R<sub>2</sub> and R<sub>4</sub> are methyl, then R' and R'' can not be H, OH or OCH<sub>3</sub>.

15. (currently amended) The method of claim 14 wherein:

R''' ~~comprises~~ is selected from H, halogen, C(halogen)<sub>3</sub>, lower alkyl ~~or~~ and alkoxy;

R'''' ~~comprises~~ is selected from H, halogen, C(halogen)<sub>3</sub>, lower alkyl ~~or~~ and alkoxy; and

R'' ~~comprises~~ is -Y-D<sub>1</sub>-D<sub>2</sub>-T<sub>2</sub>,


~~Y comprises~~ Y is optionally present and if present ~~comprises~~ is selected from O, S, NH, N-alkyl, C=CH, C≡C, CH<sub>2</sub>, ~~CH(CH<sub>3</sub>)~~ CH(CH<sub>3</sub>), C(CH<sub>3</sub>)<sub>2</sub>, a carbocyclic ring having 4 to 6 ring members ~~or~~ and a heterocyclic ring having 4 to 6 ring members with 1 or 2 heteroatoms,

D<sub>1</sub> is optionally present and if present ~~comprises~~ is alkyl,

D<sub>2</sub> ~~comprises~~ is selected from H, an alkyl, NH, N-alkyl, O-alkyl, S-alkyl, a carbocyclic ring, a bicyclic ring, a tricyclic ring, an aromatic ring ~~or~~ and a heteroaromatic ring,

T<sub>2</sub> is optionally present and if present ~~comprises~~ is selected from an aromatic ring, a heteroaromatic ring, a heterocyclic ring, H, OH, halogen ~~or~~ and a substituent group.

16. (currently amended) A method of selectively stimulating CB2 cannabinoid receptors in an individual or animal comprising administering to the individual or animal a therapeutically effective amount of at least one compound of formula I below, and physiologically acceptable salts thereof:



wherein,

the "A" ring atoms of compound formula I comprise carbon and 0 to 2 nitrogen heteroatoms;

Ar is an aromatic ring, an aromatic ring comprising at least one substituent group, a heteroaromatic ring, a heteroaromatic ring comprising 1 to 5 substituent groups, a heterocyclic ring or a heterocyclic ring comprising at least one substituent group;

R ~~comprises~~ is selected from H, OH, OCH<sub>3</sub>, alkoxy, OCH<sub>2</sub>CH<sub>2</sub>OH, alcohol, NH<sub>2</sub>, PO<sub>3</sub>H, OPO<sub>3</sub>H, OSO<sub>3</sub>H, halogen, C(halogen)<sub>3</sub>, SE<sub>1</sub>, OE<sub>1</sub> ~~or~~ and NE<sub>1</sub>E<sub>2</sub>,

E<sub>1</sub> and E<sub>2</sub> are each independently H or alkyl;

R' ~~comprises~~ is selected from H, OH, alkoxy, OCH<sub>2</sub>CH<sub>2</sub>OH, alcohol, NH<sub>2</sub>, PO<sub>3</sub>H,

OPO<sub>3</sub>H, OSO<sub>3</sub>H, halogen, C(halogen)<sub>3</sub>, SE<sub>1</sub>, OE<sub>1</sub> or and NE<sub>1</sub>E<sub>2</sub>,

E<sub>1</sub> and E<sub>2</sub> are each independently H or alkyl;

~~R''[[,]] R''' and R'''' each independently comprises is~~ Y-D<sub>1</sub>-D<sub>2</sub>-T<sub>2</sub>, ~~H, halogen, alkyl, alkoxy or a substituent group,~~

Y is optionally present and if present ~~comprises is~~ selected from O, S, NH, N-alkyl, C=CH, C≡C, CH<sub>2</sub>, ~~CH(CH<sub>3</sub>)~~ CH(CH<sub>3</sub>), C(CH<sub>3</sub>)<sub>2</sub>, a carbocyclic ring having 4 to 6 ring members or and a heterocyclic ring having 4 to 6 ring members with 1 or 2 heteroatoms,

D<sub>1</sub> is optionally present and if present ~~comprises is~~ alkyl,

D<sub>2</sub> ~~comprises is~~ selected from H, alkyl, NH, N-alkyl, O-alkyl, S-alkyl, a carbocyclic ring, a bicyclic ring, a tricyclic ring, an aromatic ring or and a heteroaromatic ring,

T<sub>2</sub> is optionally present and if present ~~comprises is~~ selected from an aromatic ring, a substituted aromatic ring, a heteroaromatic ring, a substituted heteroaromatic ring, a heterocyclic ring, a substituted heterocyclic ring, H, OH, halogen, or and a substituent group;

R''' and R'''' are each independently selected from H, halogen, alkyl, alkoxy and a substituent group;

with the proviso that,

when Ar is 4-isopropyl pyridine or 4-isopropenyl pyridine, R''' is hydrogen, and R'''' is hydrogen, then R'' can not be a straight or branched saturated alkyl having 1 to 20 carbon atoms;

when Ar is 4-isopropyl toluene or 4-isopropenyl toluene, and both R''' and R'''' are hydrogen, R'' can not be a straight or branched saturated alkyl having 1 to 20 carbon atoms;

when R'' is C(CH<sub>3</sub>)<sub>2</sub>(CH<sub>2</sub>)<sub>5</sub>CH<sub>3</sub>, R<sub>2</sub> and R<sub>4</sub> are methyl, then R' and R'' can not be H, OH or OCH<sub>3</sub>.

17. (currently amended) The method of claim 16, wherein:

R''' comprises is selected from H, halogen, C(halogen)<sub>3</sub>, lower alkyl or and alkoxy;

R''' comprises is selected from H, halogen, C(halogen)<sub>3</sub>, lower alkyl or and alkoxy; and

R'' comprises is -Y-D<sub>1</sub>-D<sub>2</sub>-T<sub>2</sub>,


~~Y comprises~~ Y is optionally present and if present comprises is selected from O, S, NH, N-alkyl, C=CH, C≡C, CH<sub>2</sub>, ~~CH(CH<sub>3</sub>)~~ CH(CH<sub>3</sub>), C(CH<sub>3</sub>)<sub>2</sub>, a carbocyclic ring having 4 to 6 ring members or and a heterocyclic ring having 4 to 6 ring members with 1 or 2 heteroatoms,

D<sub>1</sub> is optionally present and if present comprises is alkyl,

D<sub>2</sub> comprises is selected from H, an alkyl, NH, N-alkyl, O-alkyl, S-alkyl, a carbocyclic ring, a bicyclic ring, a tricyclic ring, an aromatic ring or and a heteroaromatic ring,

T<sub>2</sub> is optionally present and if present comprises is selected from an aromatic ring, a heteroaromatic ring, a heterocyclic ring, H, OH, halogen or and a substituent group.

18. (currently amended) A method of treating a condition comprising administering to an individual or animal having the condition a therapeutically effective amount of at least one compound of formula I below, and physiologically acceptable salts thereof:



wherein,

the "A" ring atoms of compound formula I comprise carbon and 0 to 2 nitrogen heteroatoms;

Ar is an aromatic ring, an aromatic ring comprising at least one substituent group,

a heteroaromatic ring, a heteroaromatic ring comprising 1 to 5 substituent groups, a heterocyclic ring or a heterocyclic ring comprising at least one substituent group;

R ~~comprises~~ is selected from H, OH, OCH<sub>3</sub>, alkoxy, OCH<sub>2</sub>CH<sub>2</sub>OH, alcohol, NH<sub>2</sub>, PO<sub>3</sub>H, OPO<sub>3</sub>H, OSO<sub>3</sub>H, halogen, C(halogen)<sub>3</sub>, SE<sub>1</sub>, OE<sub>1</sub> ~~or~~ and NE<sub>1</sub>E<sub>2</sub>,

E<sub>1</sub> and E<sub>2</sub> are each independently H or alkyl;

R' ~~comprises~~ is selected from H, OH, alkoxy, OCH<sub>2</sub>CH<sub>2</sub>OH, alcohol, NH<sub>2</sub>, PO<sub>3</sub>H, OPO<sub>3</sub>H, OSO<sub>3</sub>H, halogen, C(halogen)<sub>3</sub>, SE<sub>1</sub>, OE<sub>1</sub> ~~or~~ and NE<sub>1</sub>E<sub>2</sub>,

E<sub>1</sub> and E<sub>2</sub> are each independently H or alkyl;

R''[,.] R''' and R'''' each independently ~~comprises~~ is Y-D<sub>1</sub>-D<sub>2</sub>-T<sub>2</sub>, H, ~~halogen, alkyl, alkoxy or a substituent group,~~

Y is optionally present and if present ~~comprises~~ is selected from O, S, NH, N-alkyl, C=CH, C≡C, CH<sub>2</sub>, ~~CH(CH<sub>3</sub>)~~ CH(CH<sub>3</sub>), C(CH<sub>3</sub>)<sub>2</sub>, a carbocyclic ring having 4 to 6 ring members ~~or~~ and a heterocyclic ring having 4 to 6 ring members with 1 or 2 heteroatoms,

D<sub>1</sub> is optionally present and if present ~~comprises~~ is alkyl,

D<sub>2</sub> ~~comprises~~ is selected from H, alkyl, NH, N-alkyl, O-alkyl, S-alkyl, a carbocyclic ring, a bicyclic ring, a tricyclic ring, an aromatic ring ~~or~~ and a heteroaromatic ring,

T<sub>2</sub> is optionally present and if present ~~comprises~~ is selected from an aromatic ring, a substituted aromatic ring, a heteroaromatic ring, a substituted heteroaromatic ring, a heterocyclic ring, a substituted heterocyclic ring, H, OH, halogen, ~~or~~ and a substituent group;

R''' and R'''' are each independently selected from H, halogen, alkyl, alkoxy and a substituent group.

with the proviso that,

when Ar is 4-isopropyl pyridine or 4-isopropenyl pyridine, R''' is hydrogen, and R'''' is hydrogen, then R'' can not be a straight or branched saturated alkyl having 1 to 20 carbon atoms;

when Ar is 4-isopropyl toluene or 4-isopropenyl toluene, and both R''' and R'''' are hydrogen, R'' can not be a straight or branched saturated alkyl having 1 to 20 carbon atoms;

when R'' is C(CH<sub>3</sub>)<sub>2</sub>(CH<sub>2</sub>)<sub>5</sub>CH<sub>3</sub>, R<sub>2</sub> and R<sub>4</sub> are methyl, then R' and R'' can not be H, OH or OCH<sub>3</sub>.

19. (currently amended) The method of claim 18, wherein:

R''' comprises is selected from H, halogen, C(halogen)<sub>3</sub>, lower alkyl ~~or~~ and alkoxy;

R'''' comprises is selected from H, halogen, C(halogen)<sub>3</sub>, lower alkyl ~~or~~ and alkoxy; and

R'' comprises is -Y-D<sub>1</sub>-D<sub>2</sub>-T<sub>2</sub>,

~~Y comprises~~ Y is optionally present and if present ~~comprises~~ is selected from O, S, NH, N-alkyl, C=CH, C≡C, CH<sub>2</sub>, ~~CH(CH<sub>3</sub>)~~ CH(CH<sub>3</sub>), C(CH<sub>3</sub>)<sub>2</sub>, a carbocyclic ring having 4 to 6 ring members ~~or~~ and a heterocyclic ring having 4 to 6 ring members with 1 or 2 heteroatoms,


D<sub>1</sub> is optionally present and if present ~~comprises~~ is alkyl,

D<sub>2</sub> ~~comprises~~ is selected from H, an alkyl, NH, N-alkyl, O-alkyl, S-alkyl, a carbocyclic ring, a bicyclic ring, a tricyclic ring, an aromatic ring ~~or~~ and a heteroaromatic ring,

T<sub>2</sub> is optionally present and if present ~~comprises~~ is selected from an aromatic ring, a heteroaromatic ring, a heterocyclic ring, H, OH, halogen ~~or~~ and a substituent group.

20. (currently amended) A method of providing a physiological response in an individual or animal comprising administering to the individual or animal a therapeutically effective amount of at least one compound of formula I below, and

physiologically acceptable salts thereof:



wherein,

the "A" ring atoms of compound formula I comprise carbon and 0 to 2 nitrogen heteroatoms;

Ar is an aromatic ring, an aromatic ring comprising at least one substituent group, a heteroaromatic ring, a heteroaromatic ring comprising 1 to 5 substituent groups, a heterocyclic ring or a heterocyclic ring comprising at least one substituent group;

R ~~comprises~~ is selected from H, OH, OCH<sub>3</sub>, alkoxy, OCH<sub>2</sub>CH<sub>2</sub>OH, alcohol, NH<sub>2</sub>, PO<sub>3</sub>H, OPO<sub>3</sub>H, OSO<sub>3</sub>H, halogen, C(halogen)<sub>3</sub>, SE<sub>1</sub>, OE<sub>1</sub> ~~or~~ and NE<sub>1</sub>E<sub>2</sub>,

E<sub>1</sub> and E<sub>2</sub> are each independently H or alkyl;

R' ~~comprises~~ is selected from H, OH, alkoxy, OCH<sub>2</sub>CH<sub>2</sub>OH, alcohol, NH<sub>2</sub>, PO<sub>3</sub>H, OPO<sub>3</sub>H, OSO<sub>3</sub>H, halogen, C(halogen)<sub>3</sub>, SE<sub>1</sub>, OE<sub>1</sub> ~~or~~ and NE<sub>1</sub>E<sub>2</sub>,

E<sub>1</sub> and E<sub>2</sub> are each independently H or alkyl;

~~R''[[,]] R''' and R'''' each independently comprises~~ is ~~Y-D<sub>1</sub>-D<sub>2</sub>-T<sub>2</sub>, H, halogen, alkyl, alkoxy or a substituent group,~~

Y is optionally present and if present ~~comprises~~ is selected from O, S, NH, N-alkyl, C=CH, C≡C, CH<sub>2</sub>, CH(CH<sub>3</sub>), CH(CH<sub>3</sub>), C(CH<sub>3</sub>)<sub>2</sub>, a carbocyclic ring having 4 to 6 ring members ~~or~~ and a heterocyclic ring having 4 to 6 ring members with 1 or 2 heteroatoms,

D<sub>1</sub> is optionally present and if present ~~comprises~~ is alkyl,

D<sub>2</sub> ~~comprises~~ is selected from H, alkyl, NH, N-alkyl, O-alkyl, S-alkyl, a carbocyclic



ring, a bicyclic ring, a tricyclic ring, an aromatic ring ~~or~~ and a heteroaromatic ring,

T<sub>2</sub> is optionally present and if present ~~comprises~~ is selected from an aromatic ring, a substituted aromatic ring, a heteroaromatic ring, a substituted heteroaromatic ring, a heterocyclic ring, a substituted heterocyclic ring, H, OH, halogen, ~~or~~ and a substituent group;

R''' and R'''' are each independently selected from H, halogen, alkyl, alkoxy and a substituent group.

with the proviso that,

when Ar is 4-isopropyl pyridine or 4-isopropenyl pyridine, R''' is hydrogen, and R'''' is hydrogen, then R'' can not be a straight or branched saturated alkyl having 1 to 20 carbon atoms;

when Ar is 4-isopropyl toluene or 4-isopropenyl toluene, and both R''' and R'''' are hydrogen, R'' can not be a straight or branched saturated alkyl having 1 to 20 carbon atoms;

when R'' is C(CH<sub>3</sub>)<sub>2</sub>(CH<sub>2</sub>)<sub>5</sub>CH<sub>3</sub>, R<sub>2</sub> and R<sub>4</sub> are methyl, then R' and R'' can not be H, OH or OCH<sub>3</sub>.

21. (currently amended) The method of claim 20, wherein:

R''' ~~comprises~~ is selected from H, halogen, C(halogen)<sub>3</sub>, lower alkyl ~~or~~ and alkoxy;

R'''' ~~comprises~~ is selected from H, halogen, C(halogen)<sub>3</sub>, lower alkyl ~~or~~ and alkoxy; and

R'' ~~comprises~~ is -Y-D<sub>1</sub>-D<sub>2</sub>-T<sub>2</sub>,

~~Y comprises~~ Y is optionally present and if present ~~comprises~~ is selected from O, S, NH, N-alkyl, C=CH, C≡C, CH<sub>2</sub>, ~~CH(CH<sub>3</sub>)~~ CH(CH<sub>3</sub>), C(CH<sub>3</sub>)<sub>2</sub>, a carbocyclic ring having 4 to 6 ring members ~~or~~ and a heterocyclic ring having 4 to 6 ring members with 1 or 2 heteroatoms,


D<sub>1</sub> is optionally present and if present ~~comprises~~ is alkyl,

D<sub>2</sub> ~~comprises~~ is selected from H, an alkyl, NH, N-alkyl, O-alkyl, S-alkyl, a carbocyclic ring, a bicyclic ring, a tricyclic ring, an aromatic ring ~~or~~ and a

heteroaromatic ring,

T<sub>2</sub> is optionally present and if present ~~comprises~~ is selected from an aromatic ring, a heteroaromatic ring, a heterocyclic ring, H, OH, halogen ~~or~~ and a substituent group.

22. (currently amended) A method of treating a condition selected from central and peripheral pain, neuropathy, neurodegenerative diseases including multiple sclerosis, Parkinson's disease, Huntington's chorea, Alzheimer's disease; mental disorders such as schizophrenia and depression, endotoxic shock, hypotensive shock; or of modulating appetite; or of modulating the immune system; or of reducing fertility; or of treating diseases associated with motor function such as Tourette's syndrome; or of treating inflammation; or of providing neuroprotection; or of suppressing memory; or of producing peripheral vasodilation; or of treating epilepsy, glaucoma, nausea associated with cancer chemotherapy or nausea associated with Aids wasting syndrome comprising administering to an individual or animal having the condition a therapeutically effective amount of at least one compound ~~at least one compound~~ of formula I below, and physiologically acceptable salts thereof:



wherein,

the "A" ring atoms of compound formula I comprise carbon and 0 to 2 nitrogen heteroatoms;

Ar is an aromatic ring, an aromatic ring comprising at least one substituent group, a heteroaromatic ring, a heteroaromatic ring comprising 1 to 5 substituent groups, a heterocyclic ring or a heterocyclic ring comprising at least one substituent group;

R ~~comprises~~ is selected from H, OH, OCH<sub>3</sub>, alkoxy, OCH<sub>2</sub>CH<sub>2</sub>OH, alcohol, NH<sub>2</sub>,

$\text{PO}_3\text{H}$ ,  $\text{OPO}_3\text{H}$ ,  $\text{OSO}_3\text{H}$ , halogen,  $\text{C}(\text{halogen})_3$ ,  $\text{SE}_1$ ,  $\text{OE}_1$  or and  $\text{NE}_1\text{E}_2$ ,

$\text{E}_1$  and  $\text{E}_2$  are each independently H or alkyl;

$\text{R}'$  ~~comprises~~ is selected from H, OH, alkoxy,  $\text{OCH}_2\text{CH}_2\text{OH}$ , alcohol,  $\text{NH}_2$ ,  $\text{PO}_3\text{H}$ ,  $\text{OPO}_3\text{H}$ ,  $\text{OSO}_3\text{H}$ , halogen,  $\text{C}(\text{halogen})_3$ ,  $\text{SE}_1$ ,  $\text{OE}_1$  or and  $\text{NE}_1\text{E}_2$ ,

$\text{E}_1$  and  $\text{E}_2$  are each independently H or alkyl;

$\text{R}''$  ~~[[,]]~~  $\text{R}'''$  and  $\text{R}''''$  each independently ~~comprises~~ is  $\text{Y-D}_1\text{-D}_2\text{-T}_2$ , H, ~~halogen, alkyl,~~  
~~alkoxy or a substituent group,~~

Y is optionally present and if present ~~comprises~~ is selected from O, S, NH, N-alkyl,  $\text{C}=\text{CH}$ ,  $\text{C}\equiv\text{C}$ ,  $\text{CH}_2$ ,  $\text{CH}(\text{CH}_3)$ ,  $\text{CH}(\text{CH}_3)$ ,  $\text{C}(\text{CH}_3)_2$ , a carbocyclic ring having 4 to 6 ring members or and a heterocyclic ring having 4 to 6 ring members with 1 or 2 heteroatoms,

$\text{D}_1$  is optionally present and if present ~~comprises~~ is alkyl,

$\text{D}_2$  ~~comprises~~ is selected from H, alkyl, NH, N-alkyl, O-alkyl, S-alkyl, a carbocyclic ring, a bicyclic ring, a tricyclic ring, an aromatic ring or and a heteroaromatic ring,

$\text{T}_2$  is optionally present and if present ~~comprises~~ is selected from an aromatic ring, a substituted aromatic ring, a heteroaromatic ring, a substituted heteroaromatic ring, a heterocyclic ring, a substituted heterocyclic ring, H, OH, halogen, or and a substituent group;

$\text{R}'''$  and  $\text{R}''''$  are each independently selected from H, halogen, alkyl, alkoxy and a substituent group,

with the proviso that,

when Ar is 4-isopropyl pyridine or 4-isopropenyl pyridine,  $\text{R}'''$  is hydrogen, and  $\text{R}''''$  is hydrogen, then  $\text{R}''$  can not be a straight or branched saturated alkyl having 1 to 20 carbon atoms;

when Ar is 4-isopropyl toluene or 4-isopropenyl toluene, and both  $\text{R}'''$  and  $\text{R}''''$  are hydrogen,  $\text{R}''$  can not be a straight or branched saturated alkyl having 1 to 20 carbon atoms;

when R'' is C(CH<sub>3</sub>)<sub>2</sub>(CH<sub>2</sub>)<sub>5</sub>CH<sub>3</sub>, R<sub>2</sub> and R<sub>4</sub> are methyl, then R' and R'' can not be H, OH or OCH<sub>3</sub>.

23. (currently amended) The method of claim 22, wherein:

R''' ~~comprises~~ is selected from H, halogen, C(halogen)<sub>3</sub>, lower alkyl ~~or~~ and alkoxy;

R'''' ~~comprises~~ is selected from H, halogen, C(halogen)<sub>3</sub>, lower alkyl ~~or~~ and alkoxy; and

R'' ~~comprises~~ is -Y-D<sub>1</sub>-D<sub>2</sub>-T<sub>2</sub>,

~~Y comprises~~ Y is optionally present and if present ~~comprises~~ is selected from O, S, NH, N-alkyl, C=CH, C≡C, CH<sub>2</sub>, ~~CH(CH<sub>3</sub>)~~ CH(CH<sub>3</sub>), C(CH<sub>3</sub>)<sub>2</sub>, a carbocyclic ring having 4 to 6 ring members ~~or~~ and a heterocyclic ring having 4 to 6 ring members with 1 or 2 heteroatoms,

D<sub>1</sub> is optionally present and if present ~~comprises~~ is alkyl,

D<sub>2</sub> ~~comprises~~ is selected from H, an alkyl, NH, N-alkyl, O-alkyl, S-alkyl, a carbocyclic ring, a bicyclic ring, a tricyclic ring, an aromatic ring ~~or~~ and a heteroaromatic ring,

T<sub>2</sub> is optionally present and if present ~~comprises~~ is selected from an aromatic ring, a heteroaromatic ring, a heterocyclic ring, H, OH, halogen ~~or~~ and a substituent group.